

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE 1-25-07	3. REPORT TYPE AND DATES COVERED Final 01/01/06 - 12/31/06		
4. TITLE AND SUBTITLE Predictability and Dynamics of Geophysical Fluid Flows - GFA Extension		5. FUNDING NUMBERS N00014-06-1-1369		
6. AUTHOR(S) Roger M. Samelson				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Oregon State University Corvallis, OR 97331		8. PERFORMING ORGANIZATION REPORT NUMBER N02100		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) ONR		10. SPONSORING/MONITORING AGENCY REPORT NUMBER		
11. SUPPLEMENTARY NOTES DISTRIBUTION STATEMENT A Approved for Public Release Distribution Unlimited				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Unlimited Public Access		12b. DISTRIBUTION CODE		
13. ABSTRACT (Maximum 200 words) Research under this grant, an extension of Grant #N00014-98-1-0813 that supported the completion of the Ph.D. thesis of Christopher L. Wolfe, focused primarily on computations of unstable nonlinear periodic solutions, time-dependent normal modes (Floquet vectors), and singular vectors in a two-layer quasi-geostrophic channel model. The model was studied in a strongly nonlinear regime, in which small disturbances to an unstable, steady, zonal, baroclinic shear flow grow to finite amplitude and continue to vacillate irregularly for arbitrarily long times. The computation of time-dependent, normal-mode disturbances to unstable, nonlinear, time-periodic basic flows in a high-dimensional geophysical fluid model opens a new perspective on the analysis of disturbance growth in time-dependent flows, and on the closely related problem of error growth in predictive models of time-dependent flows.				
14. SUBJECT TERMS		15. NUMBER OF PAGES		
		16. PRICE CODE		
17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT	

FINAL TECHNICAL REPORT

ONR GRANT #N00014-06-1-1369

PI: Roger M. Samelson

Title: Predictability and Dynamics of Geophysical Fluid Flows – GRA Extension

Research under this grant, an extension of Grant #N00014-98-1-0813 that supported the completion of the Ph.D. thesis of Christopher L. Wolfe, focused primarily on computations of unstable nonlinear periodic solutions, time-dependent normal modes (Floquet vectors), and singular vectors in a two-layer quasi-geostrophic channel model. The model was studied in a strongly nonlinear regime, in which small disturbances to an unstable, steady, zonal, baroclinic shear flow grow to finite amplitude and continue to vacillate irregularly for arbitrarily long times. The computation of time-dependent, normal-mode disturbances to unstable, nonlinear, time-periodic basic flows in a high-dimensional geophysical fluid model opens a new perspective on the analysis of disturbance growth in time-dependent flows, and on the closely related problem of error growth in predictive models of time-dependent flows.

The results of this one-year extension confirm and extend the results reported for the previous Grant #N00014-98-1-0813. In the strongly nonlinear regime, unstable periodic solutions were identified and computed using an efficient Newton-Picard method. The complete Floquet and singular vector spectra were computed for strongly nonlinear cycles and the dynamical structure of the resulting modes were analyzed. The Floquet vectors fall into two classes with direct physical interpretations: wave dynamical (WD) modes and damped-advective (DA) modes. Singular vectors were computed and analyzed in terms of the two classes of normal-mode solutions, and showed a similar dynamical splitting. A new, efficient method was developed for the recovery of Lyapunov vectors from singular vectors, which promises the possibility of computing Lyapunov vectors directly for the first time for very large systems such as numerical weather prediction models.

LIST OF PUBLICATIONS

- Wolfe, C. L., and Samelson, R. M., 2006. Normal mode analysis of a baroclinic wave-mean oscillation. *Journal of the Atmospheric Sciences*, 63, 2795-2812.
- Wolfe, C. L., and R. M. Samelson, 2007. An efficient method for recovering Lyapunov vectors from singular vectors. *Tellus*, accepted.
- Wolfe, C. L., and R. M. Samelson. Singular vectors and time-dependent normal modes of a baroclinic wave-mean oscillation. *Journal of the Atmospheric Sciences*, submitted.

DISTRIBUTION STATEMENT A
Approved for Public Release
Distribution Unlimited

20070131182